Percent of women, ages 18 through 44, with a preventive medical visit in the past year

GOAL	To increase the percent of women who have an annual preventive medical visit.	
DEFINITION	Numerator: Number of women, ages 18 through 44, who had a preventive medical visit in the past year Denominator: Number of women, ages 18 through 44	
	<b>Units:</b> 100	Text: Percent
HEALTHY PEOPLE 2020 OBJECTIVE		
	Related to Access to Health Services Objective 7.0: Increase the proportion appropriate clinical preventive service	n of persons who receive
DATA SOURCES and DATA ISSUES	Behavioral Risk Factor Surveillance System (BRFSS)	
MCH POPULATION DOMAIN	Women/Maternal Health	
SIGNIFICANCE	A well-woman or preconception visit opportunity to receive recommended services, including screening, counse which can lead to appropriate identifi prevention of disease to optimize the between, and beyond potential pregr screening and management of chron diabetes, and counseling to achieve smoking cessation, can be advanced to promote women's health prior to a and improve subsequent maternal ar The annual well-woman visit is recon College of Obstetrics and Gynecolog http://www.acog.org/Resources-And- Opinions/Committee-on-Gynecologic Visit	clinical preventive eling, and immunizations, ication, treatment, and e health of women before, hancies. For example, hic conditions such as a healthy weight and d within a well woman visit and between pregnancies and perinatal outcomes. hmended by the American hists (ACOG). -Publications/Committee-

PERFORMANCE MEASURE 14	14.1 Percent of women who smoke during pregnancy and 14.2 Percent of children, ages 0 through 17, who live in households where someone smokes	
GOAL	<ul><li>14.1 To decrease the number of women who smoke during pregnancy and</li><li>14.2 To decrease the number of households where someone smokes.</li></ul>	
DEFINITION	<ul> <li>Numerator:</li> <li>14.1 Number of women who report smoking during pregnancy</li> <li>14.2 Number of children, ages 0 through 17, who live in households where there is household member who smokes</li> </ul>	
	<b>Denominator:</b> 14.1 Number of live births 14.2 Number of children, ages 0 through 17	
	Units: 100 Text: Percent	
HEALTHY PEOPLE 2020 OBJECTIVE	Related to Tobacco Use (TU) Objective 6: Increase smoking cessation during pregnancy (Target: 30.0%) and related to Tobacco Use (TU) Objective 11.1: Reduce the proportion of children aged 3 to 11 years exposed to secondhand smoke. (Baseline: 52.2%, Target 47%)	
DATA SOURCES and DATA ISSUES	14.1 National Vital Statistics System (NVSS) 14.2 National Survey of Children's Health (NSCH)	
MCH POPULATION DOMAIN	Women/Maternal Health, Child Health, and/or Adolescent Health	
SIGNIFICANCE	Women who smoke during pregnancy are more likely to experience a fetal death or deliver a low birth weight baby. Adverse effects of parental smoking on children have been a clinical and public health concern for decades. Children have an increased frequency of ear infections; acute respiratory illnesses and related hospital admissions during infancy; severe asthma and asthma-related problems; lower respiratory tract infections; and SIDS.	
	The Health Consequences of Smoking—50 Years of Progress. A Report of the Surgeon General. Atlanta, Georgia: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2014. https://www.surgeongeneral.gov/library/reports/50-years-of- progress/index.html	